

Publications Xlab 2010

1. Olsen, D.B., Scheede-Bergdahl, C., Reving, D., Boushel, R., Dela, F. The effect of rowing on endothelial function and insulin action in healthy controls and in patients with type 2 diabetes. *Scandinavian Journal of Medicine and Science in Sports*. Epub ahead of print, 2010.
2. Naimi, A., Rabøl, R., Wright-Paradis, C., Rossi, A., Bourbeau, J., Dela, F., Boushel, R. Altered mitochondrial function in locomotor muscles of patients with COPD. *Clin Physiol Funct Imaging*. In press 2010.
3. Eriksen, L., Grønæk, M., Helge, J.W., Tolstrup, J.S., Curtis, T. Study Design Article. The KRAM study - the Danish Health Examination Survey 2007-2007. *Scandinavian Journal of Public Health*, in press, 2010.
4. Boushel, R., Gnaiger, E., Calbet, J.A.L., Gonzalez-Alonso, J., Wright-Paradis, C., Søndergaard, H., Ara Royo, I., Helge, J.W., Saltin, B. Excess capacity of mitochondria in the oxygen consumption cascade during exercise in humans. *Mitochondrion*, in press, 2010.
5. Sonne, M.P., Højbjerg, L., Alibegovic, A.C., Vaag, A., Stallknecht, B., Dela, F. Diminished insulin-mediated forearm blood flow and muscle glucose uptake in young men with low birth weight. *Journal of Vascular Research*, 47:139-147, 2010.
6. Rabøl, R., Larsen, S., Højberg, P.M.V., Almdal, T., Boushel, R., Haugaard, S.B., Andersen, J.L., Madsbad, S., Dela, F. Regional anatomic differences in skeletal muscle mitochondrial respiration in type 2 diabetes and obesity. *The Journal of Clinical Endocrinology & Metabolism (JCEM)*, 95(2): 857-863, 2010.
7. Alibegovic A.C., Højbjerg, L., Sonne, M., Hall, G. van, Alsted, T.J., Kiens, B., Stallknecht, B., Dela, F., Vaag, A. Increased rate of whole body lipolysis before and after nine days of bed rest in healthy young men born with low birth weight. *Am. J. Physiol. Endocrinol. Metab.*, 298: E555-E564, 2010.
8. Nordby, P., Prats, C., Kristensen, D., Ekroos, K., Forsberg, G., Andersen, J.L., Ploug, T., Dela, F., Storlien, L., Helge, J.W. Muscle ceramide content in man is higher in type I than type II fibers and not influenced by glycogen content. *European Journal of Applied Physiology*, 109:935-943, 2010.
9. Hey-Mogensen, M., Højlund K., Vind, B.F., Wang, L., Dela, F., Beck-Nielsen, H., Fernström, M., Sahlin, K. Effect of physical training on mitochondrial respiration and ROS release in skeletal muscle in obesity and type 2 diabetes. *Diabetologia*, 53:1976-1985, 2010.
10. Dela, F., Stallknecht, B. The effect of physical training on insulin secretion and action in skeletal muscle and adipose tissue of first degree relatives to type 2 diabetic patients. *American Journal of Physiology*, 299:E80-E91, 2010.

11. Kraunsøe, K., Boushel, R., Hansen, C.N., Schjerling, P., Qvortrup, K., Støckel, M., Mikines, K.J., Dela, F. Mitochondrial respiration in subcutaneous and visceral adipose tissue from patients with morbid obesity. *Journal of Physiology*, 588.12: 2023-2032, 2010.
12. Sonne, M.P., Alibegovic, A.C., Højbjerg, L., Vaag, A., Stallknecht, B., Dela, F. The effect of ten days of bed rest on metabolic and vascular insulin action. A study in individuals at risk for type 2 diabetes. *J.Appl.Physiol.*, 108:830-837, 2010.
13. Alibegovic, A.C., Sonne, M.P., Højbjerg, L., Hansen, T., Pedersen, O., Van Hall, G., Holst, J.J., Stallknecht, B., Dela, F., Vaag, A. The T-allele of TCF7L2 rs7903146 associates with a reduced compensation of insulin secretion for insulin resistance induced by 9 days of bed rest. *Diabetes*, 59: 836-843, 2010.
14. Rabøl, R., Boushel, R., Almdal, T., Hansen, C.N., Ploug, T., Haugaard, S.B., Prats, C., Madsbad, S., Dela, F. Opposite effects of pioglitazone and rosiglitazone on mitochondrial respiration in skeletal muscle of patients with type 2 diabetes. *Diabetes Obes Metab*, 12(9):806-14, 2010.
15. Alibegovic, A.C., Sonne, M.P., Højbjerg, L., Bork-Jensen, J., Jacobsen, S., Nilsson, E., Færch, K., Hiscock, N.J., Mortensen, B., Friedrichsen, M., Stallknecht, B., Dela, F., Vaag, A. Insulin resistance induced by physical inactivity is associated with multiple transcriptional changes in skeletal muscle in young men. *Am J Physiol Endocrinol Metab*, 299(5): E752-63, 2010.
16. Højbjerg, L., Sonne, M.P., Alibegovic, A.C., Dela, F., Vaag, A., Meldgaard, J.B., Christensen, K.B., Stallknecht, B. Impact of physical inactivity on subcutaneous adipose tissue metabolism in healthy young male offspring of patients with type 2 diabetes. *Diabetes*, 59(11): 2790-8, 2010.
17. Petersen, C.B., Thygesen, L.C., Helge, J.W., Grønbaek, M., Tolstrup, J.S. Time trends in physical activity in leisure time in the Danish population from 1987 to 2005. *Scand J. Public Health*, 38(2): 12-128. 2010.
18. Rosenkilde, M., Nordby, P., Nielsen, L.B., Stallknecht, B.M., Helge, J.W. Fat oxidation at rest predicts peak fat oxidation during exercise and metabolic phenotype in overweight males. *IJO*, 34(5): 871-7, 2010.
19. Kynde, I., Heitmann, B.L., Bygbjerg, I.C., Andersen, L.B., Helge, J.W. Childhood hypo-adiponectinaemia but not hyper-leptinaemia is associated with insulin insensitivity six years later. *Paediatric Diabetes*, 11(3); 195-202, 2010.
20. Helge, J.W. Arm and leg substrate utilization and muscle adaptation after prolonged low intensity training. *Acta Physiologica*, 199(4): 519-528, 2010.
21. Kynde, I., Johnsen, N.F., Bygbjerg, I.C., Wedderkop, N., Helge, J.W., Heitmann, B.L. Intake of dietary added sugar is directly associated with insulin resistance among girls but not boys. European Youth Heart Study I and II. *Public Health Nutrition*, 13(10):1669-74, 2010.

22. Ara, I., Larsen S., Stallknecht B., Guerra B., Fuentes T., Andersen J.L., Olmedillas H., Guadalupe-Grau A., Henrik Galbo, Calbet J.A.L., and Helge J.W. Normal mitochondrial function and increased fat oxidation capacity in leg and arm muscles in obese humans. *IJO*, in press, 2010.
23. Helge, J.W., Stallknecht, B., Drachmann, T., Hellgren, L.I., Jiménez-Jiménez, R., Andersen, J.L., Richelsen, B., Bruun, J.M. Improved glucose tolerance after intensive life style intervention occurs without changes in muscle ceramide or triacylglycerol in morbidly obese subjects. *Acta Phys*, in press, 2010.
24. Davidsen, P.K., Gallagher, I.J., Hartman, J.W., Tarnopolsky, M.A., Dela, F., Helge, J.W., Timmons, J.A., Philips, S.M. High responders to resistance exercise training demonstrate differential regulation of skeletal muscle microRNA expression. *J.Appl. Physiol*, in press, 2010.
25. Axelsen, L.N., Lademann, J.B., Petersen, J.S., Holstein-Rathlou, N.H., Ploug, T., Prats, C., Pedersen, H.D., Kjølbbye A.L. Cardiac and metabolic changes in long-term high fructose-fat fed rats with severe obesity and extensive intramyocardial lipid accumulation. *Am J Physiol Regul Integr Comp Physiol*, 298(6):R1560-70, 2010.
26. Jeppesen, J., Mogensen, M., Prats, C., Sahlin, K., Madsen, K., Kiens, B. FAT/CD36 is localized in sarcolemma and in vesicle-like structures in subsarcolemma regions but not in mitochondria. *J Lipid Res*, 51(6):1504-12, 2010.
27. Nielsen, J., Mogensen M., Vind, B.F., Sahlin, K., Højlund, K., Schrøder, H.D., Ordenblad, N. Increased subsarcolemmal lipids in type 2 diabetes: effect of training on localization of lipids, mitochondria and glycogen in sedentary human skeletal muscles. *Am J Physiol Endocrinol Metab*, 298(3):E706-13, 2010.
28. Heinonen, I., Kemppainen, J., Kaskinoro, K., Peltonen, J.E., Borra, R., Lindroos, M., Oikonen, V., Nuutila, P., Knuuti, J., Boushel, B. Regulation of human skeletal muscle perfusion and its heterogeneity during exercise in moderate hypoxia. *Am J Physiol Regul Integr Comp Physiol*, 299(1):R72-9, 2010.
29. Heinonen, I., Kemppainen, J., Kaskinoro, K., Peltonen, J.E., Borra, R., Lindroos, M.M., Oikonen, V., Nuutila, P., Knuuti, J., Hellsten, Y., Boushel, R., Kalliokoski, K.K. Comparison of exogenous adenosine and voluntary exercise on human skeletal muscle perfusion and perfusion heterogeneity. *J Appl Physiol*, 108(2):378-86, 2010.
30. Boushel, R. Muscle metaboreflex control of the circulation during exercise. *Acta Physiol (Oxf)* 199: 367-383, 2010
31. Fuentes, T., Ara, I., Guadalupe-Grau, A., Larsen, S., Stallknecht, B., Olmedillas, H., Santana, A., Helge, J.W., Calbet, J.A. and Guerra, B. Leptin receptor 170 kDa (OB-R170) protein expression is reduced in obese human skeletal muscle: a potential mechanism of leptin resistance. *Exp Physiol* 95: 160-171, 2010.